# **Tear duct probe**



#### What is a tear duct?

A tear duct is a tiny passage between the inner corner of the eyelid and the inside of the nose. It drains away tears and mucus.

### What is a blocked tear duct?

A blocked tear duct is when the passage becomes blocked and cannot drain well. Blocked tear ducts are common in babies and most often go away on their own. A blocked tear duct causes an overflow of tears and mucus that may cause an infection and crusting on the eyelashes.

## What is a tear duct probe?

A tear duct probe is a type of surgery. The doctor will place an object through the blocked tear duct to open it. This object does not stay in the tear duct.

A tear duct probe is needed if:

- The blocked tear duct does not go away on its own.
- There is a bad infection.
- There are repeated infections.

## How should I care for my child after surgery?

Some treatment guidelines include:

- Your child's eyes and nose may drain a small amount of blood-tinged fluid for up to 3 days. You may wipe the
  drainage away with a clean cloth.
- Keep your child home from day care or school the day of surgery. Your child needs rest.
- Give your child plenty of fluids to drink (unless the doctor tells you to limit fluids).
- Your child may go back to their normal activities and meals when they feel ready.
- Give your child medicine as advised by the doctor.
- Most children have some pain after this surgery. If you think your child is uncomfortable, you may give
  acetaminophen (Tylenol, Tempra, Panadol or a less costly store brand) for pain if prescribed by your child's
  doctor. Follow the directions on the box carefully, or ask your child's doctor how much medicine to give.
  - Do not give your child more than 5 doses of acetaminophen in a 24-hour period.
  - Do not give acetaminophen to babies younger than 3 months of age unless the doctor tells you it is OK.

## When should I call the doctor?

Call the doctor **right away** if your child has:

- Fever (temperature of 100.3°F or higher) for more than 3 days
- Fever that is 101°F or higher
- Eyelid that becomes swollen or warm
- Eyelid that has pus
- Any bright red bleeding
- Blood-tinged drainage that lasts more than 3 days

Also call if you have any questions or concerns about how your child looks or feels.

This teaching sheet contains general information only. Talk with your child's doctor or a member of your child.

In case of an urgent concern or emergency, call 911 or go to the nearest emergency department right away.