

Colonoscopy clean out

(For a child 40 pounds and over)

ATTENTION PARENT

It is **very important** to complete **all** of the clean out steps correctly to avoid having your child's colonoscopy rescheduled. Please contact your child's care team at least 48 hours ahead of your scheduled appointment if you have questions about the clean out instructions. ***If the colon is not empty, the doctor cannot do the colonoscopy.***

What supplies should I buy ahead of time?

- 1 bottle of Miralax (clearlax, glycolax or polyethylene glycol). The medicine label should say 238g (grams).
- 3 Dulcolax (bisacodyl) tablets. The medicine label should say 5mg (milligrams).
- Clear liquids. Avoid red and purple colors and juice with pulp. Some clear liquids include:
 - Clear apple or white grape juice
 - Gatorade or Powerade
 - Sprite or Ginger Ale
 - Popsicles
 - Jell-O
 - Soup broth or bouillon

When and how do I do the clean out?

On the day before the colonoscopy

Your child should **only drink clear liquids**, starting at breakfast. Do not let your child drink other liquids (including milk) or eat any food the rest of the day.



Starting after school (and no later than 4 p.m.):

- Give your child **2 Dulcolax tablets**.
- Mix the right amount of **Miralax with 64 oz** (ounces) of water, Gatorade or Powerade.
 - If your child weighs **40 to 88 pounds** – use **10 capfuls of Miralax**.
 - If your child weighs **over 88 pounds** – use **14 capfuls of Miralax**.
- Have your child drink about 10 oz of Miralax mix every 30 minutes so they finish all 64 oz **in 3 hours**.
- Once they finish the Miralax mix, give them **1 Dulcolax tablet**.
- Your child may continue to drink clear liquids until midnight. Do not let your child eat **any** food.

Your child will start having watery stools. Their stool should change from a brown color to yellow-green color. You will know the clean out is working when the color of their stool looks like the color of pickle juice.

Please note:

You may receive a call from a Surgery team member the day before the colonoscopy. They may tell you that your child cannot drink liquids or food after midnight. This is general information for patients receiving anesthesia to keep them relaxed or asleep, but your child still cannot eat. Continue to follow the clear liquid diet instructions given to you by your child's GI care team.

In case of an urgent concern or emergency, call 911 or go to the nearest emergency department right away.

Colonoscopy clean out (for a child 40 pounds and over), continued

Overnight and on the day of the colonoscopy

- Your child should not take aspirin or ibuprofen (Motrin, Advil or less costly store brand) 24 hours before the colonoscopy.
- Ask the doctor whether your child can take any medicines before the colonoscopy.
- Do not let your child eat any food.
- The time to stop drinking clear liquids will depend on their age and procedure time. Follow the clear liquid diet instructions given to you by your child's GI care team.

Why does my child need a clean out?

A clean out includes steps to empty all stool (poop) from the colon. This helps the doctor to:



- See the colon better.
- Take clear pictures
- Remove small pieces of tissue from the colon. These are called biopsies.



What if my child has formed or brown stool after the clean out?

If your child still has formed or brown stool after the clean out:

- Give 2 to 3 more capfuls of Miralax with 8 oz of water, Gatorade or Powerade, **and**
- Give 1 more Dulcolax tablet.



When should I call the doctor?

Call 404-785-5437 if:

- You have questions ahead of time about how to do the clean out.
- Your child is unable to complete the clean out for any reason.
- You have any other questions or concerns.

Where can I find more information?

Visit these websites to learn more about colonoscopies:

- Children's Gastroenterology Program:
choa.org/medical-services/gastroenterology
- GI Kids Colonoscopy Guide for Parents, Children and Teens:
gikids.org/tests-procedures/colonoscopy-for-kids

In case of an urgent concern or emergency, call 911 or go to the nearest emergency department right away.

Colonoscopy clean out (for a child 40 pounds and over), continued

How can I comfort my child during the clean out?

Your child may feel some discomfort during the clean out. Listed below are tips to help your child. You can also learn more at strong4life.com/en/emotional-wellness/coping/healthy-coping-strategies-for-kids.

Possible symptoms	How to help
Abdominal (belly) cramps	<ul style="list-style-type: none">• Put a warm pad on their belly.• Let them take a warm bath• Give them acetaminophen (Tylenol or less costly store brand).
Nausea (feels sick to their stomach or like throwing up)	<ul style="list-style-type: none">• Offer distractions and a comfortable environment.• Practice deep breathing with them.<ul style="list-style-type: none">– Breathe in through your nose and feel the stomach fill with air.– Hold your breath for a few seconds.– Breathe out slowly through your mouth.– Repeat several times.• Let them take a break from drinking the Miralax for 20 to 30 minutes.
Sore bottom	<ul style="list-style-type: none">• Use wet wipes instead of toilet paper.• Put diaper cream on their bottom.
Trouble drinking Miralax	<ul style="list-style-type: none">• Miralax can slowly thicken after mixing it with liquid. You can split the powder into smaller equal parts to avoid this.<ul style="list-style-type: none">– If using 3 parts, mix each with about 20 oz of clear liquid. Have them drink 1 part at a time. Then mix the next part after they finish.– You can use a different liquid for each part.– Add ice if preferred.• Turn drinking the Miralax into a game – who can finish 10 oz every 30 minutes the fastest. You can drink something that looks like what your child is drinking.• Encourage them to keep drinking.

Children's Healthcare of Atlanta has not reviewed all of the sites listed as resources and does not make any representations regarding their content or accuracy. Children's Healthcare of Atlanta does not recommend or endorse any particular products, services or the content or use of any third party websites, or make any determination that such products, services or websites are necessary or appropriate for you or for the use in rendering care to patients. Children's Healthcare of Atlanta is not responsible for the content of any of the above-referenced sites or any sites linked to these Sites. Use of the links provided on this or other sites is at your sole risk.

This teaching sheet contains general information only. Talk with your child's doctor or a member of your child's healthcare team about specific care of your child.

In case of an urgent concern or emergency, call 911 or go to the nearest emergency department right away.